§520.310

- (2) Nos. 000115, 000859, 055529, and 062250 for use of product described in paragraph (a)(1) as in paragraph (d) of this section.
 - (c) [Reserved]
- (d) Conditions of use in dogs(1) Amount. 2 mg per pound (/lb) of body weight once daily or 1 mg/lb twice daily. For the control of postoperative pain, administer approximately 2 hours before the procedure.
- (2) Indications for use. For the relief of pain and inflammation associated with osteoarthritis and for the control of postoperative pain associated with soft tissue and orthopedic surgeries.
- (3) *Limitations*. Federal Law restricts this drug to use by or on the order of a licensed veterinarian.
- [61 FR 66581, Dec. 18, 1996, as amended at 64 FR 32181, June 16, 1999; 66 FR 63165, Dec. 5, 2001; 67 FR 6866, Feb. 14, 2002; 67 FR 65038, Oct. 23, 2002; 67 FR 65697, Oct. 28, 2002; 70 FR 30626, May 27, 2005; 71 FR 51995, Sept. 1, 2006; 72 FR 68478, Dec. 5, 2007; 74 FR 21768, May 11, 2009; 78 FR 52853, Aug. 27, 2013; 78 FR 66264, Nov. 5, 2013]

§ 520.310 Caramiphen ethanedisulfonate and ammonium chloride tablets.

- (a) Specifications. Each tablet contains 10 milligrams of 5st caramiphen ethanedisulfonate and 80 milligrams of ammonium chloride.¹
- (b) *Sponsor*. See No. 000856 in §510.600(c) of this chapter.
- (c) Conditions of use in dogs(1) Amount. One tablet per 15 to 30 pounds of body weight every 4 to 6 hours.¹
- (2) Indications for use. For relief of cough.1

[43 FR 55385, Nov. 28, 1978]

§520.312 Carnidazole tablets.

- (a) *Specifications*. Each tablet contains 10 milligrams of carnidazole.
- (b) *Sponsor*. See 053923 in §510.600(c) of this chapter.
- (c) Conditions of use(1) Amount. Adult pigeons: 1 tablet (10 milligrams); newly weaned pigeons: ½ tablet (5 milligrams).

- (2) Indications for use. For treating trichomoniasis (canker) in ornamental and homing pigeons.
- (3) Limitations. Not for use in pigeons intended for human food. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism or when severely ill birds do not respond to treatment.

[54 FR 32336, Aug. 7, 1989]

§520.314 Cefadroxil.

- (a) *Specifications*.(1) Each tablet contains 50, 100, or 200 milligrams (mg) or 1 gram of cefadroxil.
- (2) Each milliliter of suspension constituted from powder contains 50 mg of cefadroxil.
- (b) *Sponsor*. See No. 000010 in §510.600(c) of this chapter.
- (c) Conditions of use in dogs and cats—(1) Amount—(i) Dogs. Administer 10 mg per pound (/lb) body weight twice daily orally.
- (ii) Cats. Administer 10 mg/lb body weight once daily orally.
- (2) Indications for use—(i) Dogs. For the treatment of skin and soft tissue infections including cellulitis, pyoderma, dermatitis, wound infections, and abscesses due to susceptible strains of Staphylococcus aureus. For the treatment of genitourinary tract infections (cystitis) due to susceptible strains of Escherichia coli, Proteus mirabilis, and S. aureus.
- (ii) Cats. For the treatment of skin and soft tissue infections including abscesses, wound infections, cellulitis, and dermatitis caused by susceptible strains of Pasteurella multocida, S. aureus, Staphylococcus epidermidis, and Streptococcus spp.
- (3) Limitations. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[75 FR 10165, Mar. 5, 2010]

§520.370 Cefpodoxime tablets.

- (a) Specifications. Each tablet contains cefpodoxime proxetil equivalent to 100 or 200 milligrams (mg) cefpodoxime.
- (b) *Sponsors*. See Nos. 000009 and 026637 in §510.600(c) of this chapter.
- (c) Conditions of use in dogs(1) Amount. 5 to 10 mg per kilogram (2.3 to 4.5 mg per pound) body weight daily for 5 to 7

¹These conditions are NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as specified by §514.111 of this chapter, but may require bioequivalency and safety information.